

Detection of Extended Spectrum Beta Lactamases among Gram Negative Bacilli Isolated From Surgical Site Infection in a Tertiary Care Hospital

Chopade Apeksha^{1*}, Ghatole Mangala², Kashetty Virendra³

^{1*}Assistant Professor, Department of Microbiology, Dr V.M.Govt. Medical College, Solapur, India.

²Professor & Head, Department of Microbiology, Ashwini Rural Medical College, Kumbhari, Solapur, India.

³Professor, Department of Microbiology, Ashwini Rural Medical College, Kumbhari, Solapur, India.

Corresponding Author: Dr. Apeksha Kundan Chopade, Assistant Professor, Department of Microbiology, Dr V.M.Govt. Medical College, Solapur, India.

Email: ^{1*}dr.apekshachopade@gmail.com

Received: 15.02.25, Revised: 19.03.25 Accepted: 07.04.25

ABSTRACT

Background: Surgical site infection (SSI) is a common complication that contribute to significant morbidity and mortality. The common organisms encountered in SSI are Staphylococcus aureus, Enterococci, E.Coli, Klebsiella spp and Pseudomonas. Knowledge of antibiotic susceptibility and drug resistant pattern will be of great help to treating physician and patient. This study is carried out to determine the prevalence of ESBL producing gram negative bacteria associated with SSI. **Material & Methods:** This is a prospective study carried out in a tertiary care hospital for a period of 2.5 years. 248 cases were diagnosed as having surgical site infection. Pus samples were processed for culture and antibiotic sensitivity. ESBL detection among gram negative bacteria was done by Predictor disk method and AmpC detection was done by AmpC disk test **Result:** Out of the 248 pus samples 221 were culture positive. Gram negative isolates 171 (69.35%) were predominantly isolated as compared to gram positive 86 (34.68%). Among the gram negative isolates 133 (77.78%) were ESBL producer of which 33.33% were pure ESBL producer, 27.48% isolates shown both ESBL & AmpC production while 16.95% isolates were pure AmpC producer. These isolates were sensitivity to Imipenem & Amikacin. **Conclusion:** Present study indicated high prevalence of ESBL producing bacteria associated with SSI. To prevent such type of infections change of antibiotic prophylaxis, screening for ESBL colonization, better infection control practices should be considered.

INTRODUCTION

Surgical site infection is a common healthcare associated infection. Even with advancement of infection prevention and control practices, better surgical techniques, antimicrobial prophylaxis, better wound care, surgical site infection still account for a major cause of prolonged hospital stays, morbidity and mortality. Surgical site infections are infections at the site of surgery within 30 or 90 days of the operation, depending on the procedure. Procedures warranting a 90-day surveillance period for the development of a surgical site infection include breast surgery, cardiac surgery, coronary artery bypass graft with both chest and donor site incisions, coronary artery bypass graft with chest incision only, craniotomy, spinal fusion, open reduction of fracture, herniorrhaphy, hip prosthesis, knee prosthesis, pacemaker surgery, peripheral vascular bypass surgery, and ventricular shunt placement.¹

Source of SSIs include the patient's own normal flora or organisms present in the hospital

environment that are introduced into the patient by medical procedures, specific underlying disease, trauma or burns which may cause a mucosal or skin surface interruption. Factors which promote SSIs include length of hospital stay, obesity, Diabetes mellitus, smoking etc. The development of a surgical site infection depends on the complex interplay of many factors. Most postoperative wounds are endogenous.²

Common organisms associated with SSI are mostly gram positive bacteria such as Staphylococcus aureus, coagulase negative Staphylococcus, Enterococcus species^{3,4} Many reports are associated with rising trend of gram negative bacilli mainly enterobacteriaceae.⁵

Isolation of gram negative bacteria may be associated with intestinal perforation, prolonged hospitalization. Many of these gram negative bacteria are ESBL producing. Risk factor for the development of such type of infections are prolonged hospital stay, inadvert

antibiotic use, previous colonization with such organism⁶

The beta lactam antibiotics are the commonly prescribed antibiotics especially given as an empirical therapy in intensive care units. Most of these beta lactam antibiotics had developed resistance by producing beta lactamase and it's a major concern in these days. The Extended spectrum beta lactamase (ESBL) are plasmid mediated enzymes that hydrolyze the oxyimino beta-lactam (3rd generation cephalosporins) and monobactams (aztreonam) but has no effect on cephamycins (cefoxitin and cefotetan).⁷ Infections caused by them are multidrug resistant and are very difficult to treat and led to use of an expensive broad spectrum antibiotics. Identification of the causative organism and its antibiotic sensitivity pattern along with ESBL detection can help in timely management of surgical site infections.

Aims & Objectives

1. To isolate and identify gram negative bacteria from Surgical site infection
2. To detect the ESBL production by phenotypic methods.

MATERIAL AND METHODS

The present study was a prospective study carried out in the Department of Microbiology

in a tertiary care hospital. The patients who developed surgical site infection were selected from General surgery, Obstetrics & Gynaecology and Orthopaedic wards

The sample size was calculated using the formula⁸

$$N = Z^2 (1 - \alpha/2) p (1 - P)$$

D^2

P = with anticipated population proportion-10%

D = Allowed error 5%.

Confidence level 95%

Patients developing stitch abscess, episiotomy wound infections were excluded from study.

After obtaining verbal consent, pus samples were collected from 248 patients suffering from surgical site infection. Swabs were collected from deeper aspect of wound and transported immediately to the laboratory. Samples were processed for microscopy and cultured on Blood agar & MacConkeys agar. Isolates were identified based on the colony characteristic, biochemical reactions as per standard microbiological protocols^{9,10}

antibiotic susceptibility testing was done on Muller Hinton agar as per CLSI guidelines.^{11,12}

ESBL detection was done by predictor disk method.¹³ Antibiotic disk were placed as shown in figure.




Figure 1

ESBL producers showed resistance to cefotaxime, ceftazidime, ceftriaxone and susceptible to cefoxitin. Increase in zone diameter by 5 mm with addition of inhibitor. Zone of Ceftazidime (30µm) and clavulanic acid (10 µm) > zone of Ceftazidime by 5mm.

Screening of Amp C was done if isolate is showing blunting of zone of any cephalosporin towards Imipenem and no increase in zone with addition of inhibitor. Confirmation of AmpC is done with AmpC disk test¹⁴ the combination with clavulanic acid bringing the susceptibility back confirms the ESBL production. A standard ESBL producing organism is usually susceptible

to cefoxitin. If there is an improvement with clavulanic acid, but not to the completely susceptible range, it would suggest either a depressed AmpC + ESBL or could also suggest the presence of an ESBL with several other (non-AmpC) enzymes. High level AmpC production has minimal effect on activity of cefepime making this drug more reliable for ESBL detection in presence of AmpC. *Escherichia coli* ATCC 25922, *Klebsiella pneumoniae* ATCC 700603 for ESBL, *Staphylococcus aureus* ATCC 25923, were used as control strains.

RESULT

248 cases were diagnosed as having surgical site infection. Among them 79 (31.85%) were male and 169 (68.15%) were female. Predominantly cases were from Obstetrics & gynecology ward. Common surgery infected was LSCS

Common age group affected was 21-30 yrs. gram negative isolates were predominant as shown in the graph Staphylococcus aureus was the predominant organism isolated. Other organisms isolated as shown in chart




Figure 2




Figure 3

All the gram-negative isolates in the study exhibited maximum sensitivity to Carbapenems Imipenem, (93%) Amikacin (71%),

Gentamicin (66%) and lesser sensitivity to 3rd generation cephalosporin's (14.15%) ESBL detection were observed as shown in the following table

Table no. 1 Incidence of ESBL, AmpC, production in Gram negative Isolates (n= 171)

| Organisms | ESBL | ESBL + AmpC | AmpC |
|---------------------------------|-------------|-------------|------------|
| E.coli N=50 | 18 (36%) | 13 (26%) | 8 (16%) |
| Acinetobacter baumannii N=46 | 6 (13.4%) | 21 (45.65%) | 5 (10.86%) |
| Klebsiella spp. N=42 | 19 (45.23%) | 9 (21.42%) | 7 (16.67%) |
| Citrobacter spp. N=12 | 5 (41.67%) | 3 (25%) | 3 (25%) |
| Pseudomonas aeruginosa N=9 | 4 (44.44%) | 0 (0) | 0 (0) |
| Proteus spp. N=8 | 5 (62.50%) | 0 (0) | 3 (37.5%) |
| Enterobacter spp. | 0 (0) | 1 (25%) | 3 (75%) |

| | | | |
|----------------|------------|------------|------------|
| N=4 | | | |
| Total N=171 | 57(33.33%) | 47(27.48%) | 29(16.95%) |

ESBL producing isolates n= 57 shown maximum susceptibility to Imipenem (100%) followed by Amikacin (75.95%), Gentamicin (47.36%) Ciprofloxacin (45.61%), tetracyclin(24.56%), Trimethoprim sulphamethaxazole(24.56%) Pure AmpC producing isolates n= 29 shown maximum susceptibility to Imipenem (100%) followed by Amikacin (57.60%), Gentamicin (34.62%) Ciprofloxacin (26.92%), Tetracyclin(15.38%), Trimethoprim sulphamethaxazole(3.85%) Both ESBL and AmpC producing isolates n= 47 shown maximum susceptibility to Imipenem (100%) followed by Amikacin (59.57%), Gentamicin (29.79%) Ciprofloxacin (8.51%), Trimethoprim sulphamethaxazole(6.38%) tetracyclin(4.25%)

DISCUSSION

Surgical site infection represents a substantial burden of diseases for patients & health services. Although the total elimination of wound infection is not possible, a reduction in the infection rate to a minimum level & spread of resistant pathogens could have significant benefits in terms of both patients comfort & resources used.

The incidence of post-operative wound infection in India ranges from 4.04- 30%.^{15,16,17} In the present study pus samples were collected from the total 248 cases of post operative wound infection. Maximum pus samples were collected from the Obstetrics & Gynaecology ward (59.27%) followed by General Surgery ward (29.03%) and Orthopaedics ward (11.69%). Predominant number of cases were operated for LSCS (30.24%). LSCS is commonly associated with the development of wound infections as various obstetric variables like prolong rupture of membrane, chorioamnionitis, and meconium increases the chances of infection.¹⁸ In this study more number of cases were from the age group of 21- 40 yrs (56.26 %) followed by 41-60 yrs (27.82%). It indicate working age group is affected. Many prospective studies on post operative wound infection pointed out that there is no evidence of relation of sex with infection of operative wound^{19,20}. In this study higher incidence of infection was found in females as compared to males. This is due to the fact that more no. of cases occurred in Obstetrics & Gynaecology ward. Mawalla et

al²¹ also showed female preponderance. Gram negative isolates (69.35%) were predominant as compared to Gram positive isolates (34.68%). This finding is in consistence with Sonawane J et al (2008)²² & Mohanty S et al (2004)²³ The most common bacterial isolate in our study was Staphylococcus aureus. Among the gram negative isolates E.coli were predominant followed by Acinetobacter baumannii, klebsiella spp. Our observations are comparable to Wassef MA et al²⁴ and Shahane V et al²⁵

Acinetobacter spp is an emerging pathogen. Recently many workers have reported it's association with wound infection and Health Care Associated Infection's. We observed higher rate of Acinetobacter spp. association with wound infection. The pattern of organisms isolated in wound types in the present study suggest the skin colonizers to be the main source of infection in clean procedures. The higher incidence of Gram negative bacilli in the clean contaminated wounds demonstrates the profound influence of endogenous contamination from the bowel and hollow muscular organs. Higher incidence of Acinetobacter spp may suggest hospital acquired infection Present study demonstrated highest number of beta lactamases producing isolates the third generation Cephalosporins are routinely used in inpatient & outpatient setting. Injudicious use result in selective pressure & emergence of drug resistance mechanism in organisms Plane ESBL production was associated with Proteus spp (62.50%), Klebsiella spp(45.23 %), Pseudomonas aeruginosa(44.44%), E.coli(36%). Our result are in comparable with other studies done with Wassef at al²⁴ & Shriyan et al.

Some workers had demonstrated ESBL production in the organisms belonging to enterobacteriaceae. There are many studies which have demonstrated ESBL production in Acinetobacter spp & Pseudomonas spp^{26,27} ESBL producing isolates were found to be susceptible to Imipenem & Amikacin which is in consistent with the observation of Sonawane J et al²². AmpC production was associated maximum in Acinetobacter baumannii, E.coli, Klebsiella spp. These isolates were found to be susceptible to Imipenem(100%) and to some extent to Amikacin (57.60%). Sarma et al²⁸ reported 14% AmpC producing E.coli from the

surgical site infection. Presence of AmpC alters the therapeutic regimen. Higher proportion of AmpC detected in our study suggest that there is a need for the continuous monitoring of detection and control of spread of these organisms. Coexistence of ESBL and AmpC production seen predominantly in *Acinetobacter baumannii* (45.65 %), *E.coli* (26%). These isolates were susceptible to Imipenem (100%) and Amikacin (57.57%). Association of AmpC with ESBL is associated with resistance to betalactam- betalactam inhibitor combination and increased amount of resistance to other classes of antibiotics.

CONCLUSION

In present study, majority of isolates from surgical site infections were *Staphylococcus aureus* 62(24.12%) followed by *E.coli* (19.34%), *Acinetobacter baumannii* 46(17.90%), *Klebsiella spp* 42(16.34%), *Coagulase negative staphylococcus* 16 (6.2%). Significant number of gram negative bacteria were identified to be ESBL Producers. ESBL production limits therapeutic option so detection of resistance pattern is important to suggest proper treatment and also help in preventing the spreads of antimicrobial resistance. Strict infection control policies should be made to prevent the spread of these resistant bacteria.

REFERENCES

- Berrios-Torres SI, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR, Reinke CE, Morgan S, Solomkin JS, Mazuski JE, Dellinger EP, Itani KMF, Berbari EF, Segreti J, Parvizi J, Blanchard J, Allen G, Kluytmans JAJW, Donlan R, Schechter WP., Healthcare Infection Control Practices Advisory Committee. Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. *JAMA Surg.* 2017 Aug 01;152(8):784-791.
- Greg J. Beilman And David L. Dunn. Surgical Infections. In Schwartz's Principles of Surgery, 9th Ed. United State of America: The McGraw-Hill Companies, Inc. 2010;113
- Giacometti A, Cirioni O, Schimizzi AM, Delprete MS, Barchiesi F, Derrico MM et al. Epidemiology and Microbiology of surgical wound infections. *J.Clin Microbiol.* 2000 feb;38(2):918-922.
- Owens CD, Stoessel K. Surgical site infections: epidemiology, microbiology and prevention. *J Hosp Infect.* 2008;70 (Suppl 2):3-10.
- Malik S, Gupta A, Singh PK, Agarwal J, Singh M. Antibigram of aerobic bacterial isolates from post- operative wound infections at a tertiary care hospital in india. *Journal of Infectious Diseases Antimicrobial Agents.* 2011;28:45-51.
- Sanjay KR, Nagendra Prasad MN, Vijaykumar GS. A study on isolation and detection of drug resistance gram negative bacilli with special importance to postoperative wound infection. *J. Microbiol. Antimicrob.* 2010; 2(6): 68-75.
- Chaudhary U, Aggarwal R. Extended spectrum -lactamases (ESBL) - An emerging threat to clinical therapeutics. *Indian J Med Microbiol.* 2004;22(2):75-80
- Lwanga SK, Lemeshao S. sample size determination in health sciences: A practical Manual. WHO publications, 25; 1991.
- Collee JG, Fraser AG, Marmion BP, Simmons A. Mackie & McCartney Practical Medical Microbiology 14th edition 2008;4(5):66-7.
- Baily FA, Sham DF, Weissfeld AS. Baily and Scott's Diagnostics Microbiology. 12th edition. Mosby Elsevier 2007:216-247.
- CLSI. Performance Standards for Antimicrobial Susceptibility Testing. 34th ed. CLSI supplement M100. Clinical and Laboratory Standards Institute; 2024
- Winn WC, Allen SD, Janda WM, Koneman EW, Procop GW, Schreckenberger PC, Woods GL editors. Antimicrobial susceptibility testing Colour atlas and textbook of diagnostic Microbiology, 6th ed. Philadelphia: Lippincott Williams and Wilkins; 2006: 947-1015
- Rodrigues C, Joshi P, Jani SH, Alphonse M, Radhakrishnan R, Mehta A. Detection of B-lactamases in gram negative clinical isolates. *Indian J Med Microbiol* 2004;22(4):247-250.
- Singhal S, Mathur T, Khan S, Upadhyay DJ, Chugh S, Gaiind R. et al. Evaluation of methods for AmpC beta-lactamase in Gram negative clinical isolates from tertiary care hospitals. *Indian J Med Microbiol* 2005; 23(2) : 120-124
- Lilani SP, Jangale N, Chowdhary A, Daver GB. Surgical site infection in clean

- and clean-contaminated cases. Indian J. Med. Microbiol. 2005; 23(4) ; 249-252
16. Suchitra Joyce B, Laxmidevi N. Surgical site infections: Assessing risk factors, outcomes and antimicrobial sensitivity patterns. Afr.JMicrobiol.Res. 2009 april; 3(4) :
17. Shriyan A, Sheetal R, Nayak N. Aerobic micro-organisms in postoperative wound infections and their antimicrobial susceptibility patterns. Journal of clinical and diagnostic research 2010 december; 4: 3392-3396
18. Amenu D, Belachew T, Araya F. Surgical site infection rate and risk factors among obstetric cases of jimma university specialised hospital, southwest Ethiopia. Ethiop J Health Sci.2011 jully; 21(2): 91-100
19. Malik S, Gupta A, Singh KP, Agarwal J, Singh M. Antibigram of aerobic bacterial isolates from postoperative wound infections at a tertiary care hospital in India. J Infect Dis Antimicrob Agents 2011; 28 : 45-51.
20. Shanthi J, Shankar T, Balagurunathan R. The scenario of post operative wound infection from a Government hospital in South India. J. Microbiol. Biotech. Res., 2012, 2 (3):396-401.
21. Mawalla B, Mshana SE, Chalya PL, Imirzalioglu C, Mathalu W. Predictors of surgical site infections among patients undergoing major surgery at Bugando medical centre in northwestern Tanzania. BMC Surg. 2011 Aug 31; 11:21
22. Sonawane J, Kamath N, Swaminathan R, Dosani K. Bacterial profile of surgical site infections and their antibiograms in a tertiary care hospital in Navi Mumbai. Bombay Hospital Journal. 2010; 52(3):358-361.
23. Mohanty S, Kapil A, Dhawan B. Bacteriological and antimicrobial susceptibility profile of soft tissue infections from Northern India. Indian Journal of Medical Sciences 2004; 58(1): 10-15
24. Wassef MA, Hussein A, Abdul Rahaman EM, El-Sherif RH. A prospective surveillance of surgical site infections: Study for efficacy of preoperative antibiotic prophylaxis. Afr. J Microbiol. Res. 2012 march; 6(12):3072-3078.
25. Shahane V, Bhawal S, Lele U. Surgical site infections: A one year prospective study in a tertiary care center. International Journal of Health Sciences, Quasim university. 2012 jan ;6(1):79-80
26. Trivedi GR, Soni ST, Vegad MM, Yadav KS. Occurrence and detection of extended spectrum B lactamase and AmpC B lactamase in clinical isolates of Pseudomonas aeruginosa and Acinetobacter baumannii by inhibitor based method. International Journal of Microbiology Research 2012; 4(8) : 299-301.
27. Sinha M, Srinivasa H, Macaden R. Antibiotic resistance profile & extended spectrum beta-lactamase (ESBL) production in Acinetobacter species. Indian J Med Res 2007; 126: 63-67.
28. Sarma JB, Bhattacharya PK, Kalita D, Rajbangshi M. Multidrug resistant Enterobacteriaceae including metallo- β -lactamase producers are predominant pathogens of healthcare-associated infections in an Indian teaching hospital. Indian J Med Microbiol 2011; 29(1) : 22-27.