

“Level of Caregivers Burden among Family Members of Mentally Ill Patients Visiting Psychiatric Outpatient Department”

Dr. Praveen Subravgoudar^{1*}, Miss. Shejal Mote², Mr. Vinay Sawant³, Miss. Shruti More⁴, Miss. Sanjana Salunkhe⁵, Miss. Sayali Kumbhar⁶, Mr. Vansh Bansode⁷, Mr. Suraj Jadhav⁸, Mr. Vinayak Yadav⁹

^{1*}M. Sc (N), Ph.D., Associate Professor, Dept. of Mental Health Nursing, D. Y. Patil College of Nursing, Kolhapur – 416003.

D. Y. Patil Education Society, (Deemed To Be University), Kolhapur, Maharashtra, India
^{2,3,4,5,6,7,8,9}Final Year B.Sc. Nursing Students, D. Y. Patil College of Nursing, Kolhapur.

D. Y. Patil Education Society, (Deemed To Be University), Kolhapur, Maharashtra, India

Corresponding Email: psubravgoudar.nsg@dypgroup.edu.in

Received: 13.06.25, Revised: 13.07.25, Accepted: 07.08.25

ABSTRACT

Background and Objectives: Mental health is a vital possession that needs to be cherished, promoted, and conserved. According to the WHO, around 450 million people worldwide suffer from mental or behavioral disorders. In India, approximately 20% of the population suffers from psychiatric or mental disorders. Caregivers play a crucial role in caring for mentally ill patients, but they often face significant emotional, physical, and financial burdens.

The researcher aims to assess the level of caregiver burden among family members of mentally ill patients visiting the psychiatric outpatient department (OPD) at Dr. D.Y. Patil Medical College Hospital and Medical Research Institute, Kolhapur. The study seeks to understand the impact of care giving on family members and identify ways to support them.

Objectives: The present study is aimed that to assess the level of caregivers burden among family members of mentally ill patients visiting psychiatric OPD at Dr. D. Y. Patil Medical College Hospital and Medical Research Institute, Kolhapur.

The objectives are as follows,

1. To assess the level of care givers burden among family member of mentally ill patients.
2. To find the association between level of care givers burden with selected demographic variables among family member of mentally ill patients.

Methods: A quantitative descriptive survey approach was used, with a non-experimental descriptive research design. A sample of 60 family members of mentally ill patients was selected using non-probability purposive sampling technique. The Zarit Caregivers Burden Scale was used to collect the data on level of caregivers burden among family member of mentally ill patients.

Result: The majority of respondents (46.66%) experienced a mild to moderate level of burden. 30% had no or minimal burden, while 20% experienced moderate to severe burden. Only 3.33% of individuals experienced a severe level of burden. Significant associations were found between caregiver burden and age, residence, and education.

Interpretation and conclusion: The study concludes that the Level of Caregiver’s Burden among Family Member of Mentally Ill Patients Visiting Psychiatric OPD at Dr. D. Y. Patil Medical College Hospital and Medical Research Institute at Kolhapur, is the majority of respondents (46.66%) experienced a mild to moderate level of burden. There is a need for healthcare professionals to assess and address caregiver burden among family members of mentally ill patients. The findings have implications for nursing practice, education, administration, and research, as well as general education in schools and colleges. Recommendations for future studies and interventions to support caregivers are also provided.

Keywords: Caregivers Burden, Family Members, Mentally Ill Patients, Psychiatric OPD, Hospital.

INTRODUCTION

They may forget what you said, but they will never forget how you made them feel

- Carl W. Buechner

-

Health especially, mental health, is one of the most important possessions of an individual and it needs to be cherished, promoted, and conserved to the maximum. Around 450 million people worldwide are suffering from

some mental or behavioral disorder according to the WHO, of which schizophrenia, bipolar disorder, depression, and alcohol use disorders are important causes for years lived with disability. According to the evidence available, in India, about 190–200/1000 populations have a psychiatric or mental disorder, this account for about 20% of the whole population. The major issues faced in India regarding mental health are lack of mental health workforce, financial aid, stigma, and caregiver burden.¹

The family plays a very vital role in the care of a mentally ill patient. A caregiver has been defined as “a family member, who has been staying with the patient for more than a year and has been closely related with the patient's daily living activities, discussions, and care of health.” Caregivers often have to sacrifice their own wants and undertake a lot of stress and are very much ignored. Caregiving drains one's emotions and hence caregivers undergo a lot of depression as compared to the general population. The WHO states caregiver burden as “the emotional, physical, financial demands and responsibilities of an individual's illness that are placed on the family members, friends, or other individuals involved with the individual outside the health-care system.” It includes taking care of personal hygiene of the patient and emotional support such as listening, counseling, giving company, and informational caring such as how to alter the living environment of the patient.¹

Burden of care can be understood by its impact and effects on caregivers. The early conceptualization of burden of care can be divided into two distinct components, objective and subjective. Objective burden of care deals with effects on finance, health, routine and leisure of the family whereas subjective burden deals with psychological and emotional impact of mental illness on family members. In addition burden of care can be precise in some upsetting notions such as shame, embarrassment, feelings of guilt, and self-blame.²

In 2004, mental disorders accounted for 13% of global burden of disease. Families who perceived a higher level of caregiver burden

are those who lived in a family with poorer functioning, worse health status and less satisfaction of social support.²

It is found that one person among four families is currently suffering from a psychiatric disorder. The families not only contribute bodily plus moral aid but also carry the adverse effects of dishonour and bias. In a family, if a family member suffers from a psychiatric illness lot of alteration and settlements have to be made by the family members to impose their role in their daily functions. Maximum time of the caregiver is consumed in taking care of the psychologically disabled patient, leading to social and economic hardships. Global burden of disease 2000 data show that neuropsychiatric disorders came out to be 30.8% of all YLDs. In fact, depression leads to maximum impairment, coming out to be 12% of all. Out of all psychiatric disorders, 6 were among top 20 sources of disability (YLDs) in the world, namely depression, alcohol abuse, bipolar disorder, schizophrenia, dementias, and headache.³

Hence the researcher wants assess the level of care givers burden among family members of mentally ill patients visiting psychiatric OPD in Dr. D.Y.Patil Medical College Hospital and Medical Research Institute at Kolhapur.

METHODS

A quantitative descriptive survey approach was used, with a non-experimental descriptive research design. A sample of 60 family members of mentally ill patients was selected using non-probability purposive sampling technique. The reliability of the tool was established and the Zarit Caregivers Burden Scale was used to collect the data on level of caregivers burden among family member of mentally ill patients.

Results.

Part I: Description of Demographic Variables of Caregivers

This part deals with distribution of participants according to their demographic characteristics. Data was analyzed using descriptive statistics and summarized in terms of percentage.

Table 1: Frequency and percentage distribution of sample according to demographic characteristics
n = 60

Variable	Frequency	Percentage
1. Age		
Less than 25 years	12	20%

25 to 35 years	25	41.66%
36 to 45 years	11	18.33%
Above 45 years	12	20%
2. Gender		
Male	29	48.33%
Female	31	51.66%
3. Religion		
Hindu	47	78.33%
Muslim	06	10%
Christian	02	3.33%
Others	05	8.33%
4. Residential area		
Urban	34	56.66%
Semi-urban	20	33.33%
Rural Area	6	10%
5. Marital Status		
Married	41	68.33%
Unmarried	18	30%
Divorced/Separated	1	1.67%
6. Type of family		
Joint	30	50%
Nuclear	27	45%
Extended	03	5%
7. Education:		
No formal education	0	0%
Primary	5	8.33%
Secondary	18	30%
Higher secondary	3	5%
Diploma	1	1.6%
Undergraduate	5	8.33%
Graduate and post graduate	28	46.66%
8. Occupation :		
Unemployment	5	8.33%
Homemaker	14	23.33%
Farmer	3	5%
Private job	27	45%
Govt. Job	4	6.6%
Business	7	11.66%
9. Monthly income of family		
<Rs 5000	02	3.33%
Rs 5001 to 10000	14	23.33%
Rs 10001 to 15000	16	26.66%
>Rs 15001	28	46.66%
10. Years of care given to the patient		
Less than 1 year	23	38.33%
1 year to 5 year	23	38.33%
5 year to 10 year	10	16.66%
More than 10 year	04	6.67%

The data presented in Table 1 shows that the majority of respondents (41.66%) belonged to the age group of 25-35 years, while 20% were less than 25 years, 20% of respondents were above 45 years. And, 18.33% were in the 36-

45 age group. Regarding gender, 51.66% of respondents were female, and 48.33% were male. In terms of religion, the majority (78.33%) were Hindu, followed by 10% who

were Muslim, 8.33% who identified with other religions, and 3.33% who were Christian.

The majority of respondents (56.66%) resided in urban areas, followed by 33.33% in semi-urban areas, and 10% in rural areas. In terms of marital status, most respondents (68.33%) were married, 30% were unmarried, and 1.67% were divorced or separated. Regarding the type of family, 50% of respondents belonged to joint families, while 45% lived in nuclear families, and 5% were part of extended families.

The majority of respondents (46.66%) had completed graduate or postgraduate education, followed by 30% who had received secondary education. A smaller percentage (8.33%) had completed primary education, while 8.33% had an undergraduate degree, 5% had a higher secondary education, and 1.6% held a diploma. None of the respondents had no formal education. In terms of occupation, 45% were employed in private jobs, 23.33% were homemakers, and 11.66%

were involved in business, 8.33% were unemployed, 6.6% held government jobs, and 5% were farmers.

The majority of respondents (46.66%) reported a monthly family income of over Rs 15,001, while 26.66% had an income between Rs 10,001 and Rs 15,000, 23.33% earned between Rs 5,001 and Rs 10,000, and only 3.33% had a family income of less than Rs 5,000. Regarding the years of care provided to the patient, 38.33% of respondents had been providing care for less than 1 year, with an equal percentage (38.33%) caring for 1 to 5 years. And, 16.66% had provided care for 5 to 10 years, and 6.67% for more than 10 years.

Part II: Assessment of the Level of Caregiver Burden among Family Members of Mentally Ill Patients.

The level of caregivers burden related to mentally ill patients among family members was assessed using Zarit caregiver burden scale.


Table 2: Frequency and percentage distribution according to the level of caregivers burden among family members of mentally ill patients. n = 60

Level of Burden	Range of score	Frequency	Percentage
No or minimal burden	0 - 20	18	30%
Mild to moderate	21 – 40	28	46.66%
Moderate to severe	41 - 60	12	20%
Severe burden	61 - 88	2	3.33%

Data in Table 2 and Figure 12 show that the majority of individuals, 46.66%, experienced a mild to moderate level of burden. 30%, had no or minimal burden, while 20% experienced

moderate to severe burden. Only 3.33% of individuals experienced a severe level of burden.

Figure 1: Distribution of the Subject According To the Level Of Caregivers Burden



Part III: Association between the Levels of Caregivers Burden with Selected Demographic Variables among Family Members.

The data reveals that the obtained Chi-square values indicate a significant association between the Level of Care givers burden with Age, Residence and Education respectively (18.381, 15.401 and 28.688 at 0.05 levels). But there is no significant association between the Level of Care givers burden and other demographic variables such as Gender, Religion, Marital Status, Type of family, Occupation, Monthly income, Years of care given.

However, the above finding reveals that there was association between the Level of Care givers burden with Age, Residence and Education respectively. So the hypothesis (H_1) was accepted.

H₁: There is a significant association between level of care givers burden with selected demographic variables among family members of mentally ill patients.

DISCUSSION

1. Demographic Characteristics

The data presented in Table 1 shows that the majority of respondents (41.66%) belonged to the age group of 25-35 years, while 20% were less than 25 years, 20% of respondents were above 45 years. And, 18.33% were in the 36-45 age group. Regarding gender, 51.66% of respondents were female, and 48.33% were male. In terms of religion, the majority (78.33%) were Hindu, followed by 10% who were Muslim, 8.33% who identified with other religions, and 3.33% who were Christian.

A study with similar finding was conducted to assess the level of family burden among caregivers of schizophrenic patients. The purposive sampling method was used to select 30 samples. The tool consisted of two sections: Section A – demographic variables and Section B – standardized questionnaire. The study result reveals that the majority of respondents (47%) were in the age group of 20-30 years and were females (70%); One-third of them were Hindu (60%); half of them were married (73.3%), were residing in rural areas (53%); and were employed (37%).²⁴

A study with similar finding was conducted to assess factors associated with the burden of family caregivers of patients with mental disorders in the psychosocial care centers, in three cities in the southwest region of Goiás state, Central Brazil. A 281 number of samples were selected by non-probability sampling

method. The study result reveals that the caregivers were mostly female and parents of the patients were married, with low education, and of low income.⁴⁵

2. Assessment of the Level of Caregivers Burden Among Family Members of Mentally Ill Patients.

The majority of individuals, 46.66%, experienced a mild to moderate level of burden. 30%, had no or minimal burden, while 20% experienced moderate to severe burden. Only 3.33% of individuals experienced a severe level of burden.

A study with similar finding was conducted to assess the burden of care among the caregivers of mentally ill patients. The quantitative approach was used. A total of 97 family members were selected consecutively from the psychiatric outpatient department (OPD) and wards of Mental hospital. The study result revealed that 40.2 % of the caregivers had mild burden, 27.8% had moderate burden, 25.8% experienced little or no burden and 6.2% experienced severe burden.²⁹

A study with contradictory findings was conducted to assess of caregiver burden in psychiatric illness patients admitted in tertiary care hospital. A sample size of 200 care givers of psychiatric patients were selected by face to face interviewing using socio-demographic and Zarit burden interview questionnaire. The study result reveals that Perception of caregiver burden found in the present study was mild in 24%, moderate in 31% and severe in 45%. Female experienced severe burden in comparison to male.²⁸

3. Association between Caregivers Burden Score with Selected Demographic Variables among Family Members Of Mentally Ill Patients.

The present study finding reveals that there was significant association between the level of care givers burden with Age, Residence and Education respectively (18.381, 15.401 and 28.688at 0.05 levels). But there is no significant association between the level of care givers burden and other demographic variables such as Gender, Religion, Marital Status, Type of family, Occupation, Monthly income, Years of care given.

A study with similar finding was conducted to assess the burden of care among the caregivers of mentally ill patients. A total of 97 family members were selected consecutively from the psychiatric outpatient department (OPD) and wards of Mental hospital. The study

result revealed that, there was significant statistical association between burden and caregiver's age ($p=0.017$), educational status ($p=0.001$), marital status ($p=0.378$), occupation ($p=0.307$), relationship with patient ($p=0.035$) and duration of care-giving ($p=0.026$). There was statistical association between burden and patient's gender ($p=0.010$), age ($p=0.40$), marital status ($p=0.50$), duration of illness ($p=0.01$) and type of illness ($p=0.09$).²⁹

A study with contradictory findings was conducted to assess the level of stress among caregivers of individuals with mental disorders attending psychiatric OPD of selected hospitals in the city. A total of 100 caregivers were selected for the study. The result reveals that, there was no significant association between age, gender, type of family, educational status, occupational status, relationship with patient diagnosis of family members, giving care to relatives, visited psychiatric OPD, no. of times admitted to psychiatric ward and significantly associated with monthly family income.⁴⁴

CONCLUSION

Majority of the caregivers participated in the study have moderate level of burden and gave free and frank responses regarding caretaker burden. The study was based on the general system model. It provides a comprehensive frame work for assessment of level of caregiversburden among family members of mentally ill patients. The research approach used is descriptive study and the samples were selected by using non-probability purposive sampling technique. Data was collected by Zarit Caregivers Burden Scale and analyzed, interpreted by applying statistical methods.

Implications of the Study

The findings of the study have implications for the nursing profession. The implications have been written under the following headings, nursing practice, nursing administration, nursing education, nursing research and general education in schools and college.

Nursing Practice

1. Assessment of Caregiver Burden: Nurses can assess caregiver burden in family members of mentally ill patients and provide support and referrals to resources as needed.

2. Individualized Care Plans: Nurses can develop individualized care plans that take into account the unique needs and circumstances of family members.
3. Support and Education: Nurses can provide support and education to family members on managing caregiver burden, stress management, and coping strategies.

Nursing Education

1. Nursing curriculum is responsible for preparing future nurses with emphasis on curative, preventive and promotive health practices.
2. Nursing educators can develop curricula that include content on caregiver burden, family-centered care, and support services for family members.
3. Nursing educators can educate students on the importance of assessing caregiver burden and providing support to family members.
4. Nursing educators can collaborate with educators from other disciplines, such as social work and psychology, to provide inter professional education on supporting family members.

Nursing Administration

1. Development of Support Services: Nursing administrators can develop support services, such as counseling, respite care, and education, to help family members manage their caregiving responsibilities and reduce their burden.
2. Staff Education and Training: Nursing administrators can provide education and training to nursing staff on the importance of assessing caregiver burden and providing support to family members.
3. Policy Development: Nursing administrators can develop policies that support family-centered care, including paid family leave, respite care, and education and training programs.
2. Resource Allocation: Nursing administrators can allocate resources, such as funding and personnel, to support the development of support services for family members.

Nursing Research

1. Longitudinal Studies: Nursing researchers can conduct longitudinal studies to examine the long-term effects of caregiving on family members of mentally ill patients.
2. Intervention Studies: Nursing researchers can design intervention studies to test the

- effectiveness of support services and other interventions in reducing caregiver burden.
3. Exploratory Studies: Nursing researchers can conduct exploratory studies to examine the experiences and needs of family members from diverse backgrounds.
 4. Instrument Development: Nursing researchers can develop instruments to measure caregiver burden and evaluate the effectiveness of support services.

General Education in Schools

1. Mental Health Education: Schools can incorporate mental health education into their curricula, including information on mental illness, treatment options, and support resources.
2. Stress Management and Coping Strategies: Schools can teach stress management and coping strategies to students, which can help them manage caregiver burden if they have a family member with a mental illness.
3. Breaking Stigmas: Schools can work to break stigmas surrounding mental illness and encourage students to seek help if they or a family member is struggling.
2. Support for Students with Family Members with Mental Illness: Schools can provide support and accommodations for students who have a family member with a mental illness, such as counseling or academic support.

General Education in College

1. Mental Health Courses: Colleges can offer courses on mental health, including the impact of mental illness on families and caregivers.
2. Caregiver Support and Resources: Colleges can provide support and resources for students who are caregivers for family members with mental illness, such as counseling or support groups.
3. Stress Management and Self-Care: Colleges can emphasize the importance of stress management and self-care for students, particularly those who are caregivers.

Limitations

1. No broad generalization could be made due to the small sample size and limited area of setting.

Recommendations

1. A similar study needs to be conducted in other healthcare settings to allow for broader generalization of caregiver burden.

2. A similar study can be conducted on a larger sample to ensure the results are representative and applicable across various populations.
3. A training program for caregivers about managing the burden of care for mentally ill patients should be periodically implemented to enhance their coping mechanisms and mental well-being.
4. A comparative study could be conducted to assess the differences in burden levels between caregivers of patients with different types of mental illnesses (e.g., schizophrenia, bipolar disorder) or between professional and non-professional caregivers.

Bibliography

1. Walke SC, Chandrasekaran V, Mayya SS. Caregiver Burden among Caregivers of Mentally Ill Individuals and Their Coping Mechanisms. *J Neurosci Rural Pract.* 2018 Apr-Jun;9(2):180-185. doi: 10.4103/jnpr.jnpr_312_17. PMID: 29725166; PMCID: PMC5912021.
2. Dhama J, Tuladhar S. Burden of Care among Caregivers of Mentally Ill Patients. *J Pharm Pract Community Med.* 2021;7(3):44-5.
3. Gupta P, Bharti P, Bathla M, Singh AH, Bhusri L. A cross-sectional study to assess the caregiver burden and the quality of life of caregivers of the patients suffering with psychiatric illness. *Ind Psychiatry J.* 2022 Jan-Jun;31(1):151-157. doi: 10.4103/ipj.ipj_228_20. Epub 2022 May 20. PMID: 35800861; PMCID: PMC9255620.
4. Cham CQ, Ibrahim N, Siau CS, Kalaman CR, Ho MC, Yahya AN, Visvalingam U, Roslan S, Abd Rahman FN, Lee KW. Caregiver Burden among Caregivers of Patients with Mental Illness: A Systematic Review and Meta-Analysis. *Healthcare (Basel).* 2022 Nov 30;10(12):2423. doi: 10.3390/healthcare10122423. PMID: 36553947; PMCID: PMC9777672.
5. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders.* Arlington: American Psychiatric Association; 2013. [Google Scholar]
6. Chou KR. Caregiver burden: a concept analysis. *J Pediatr Nurs.* 2000;15:398-399. doi:

- 10.1053/jpdn.2000.16709. [DOI] [PubMed] [Google Scholar]
7. Malhotra M. Burden among caregivers of mentally-ill patients: a review. *Int J Multidiscip Curr Res.* 2016;4:109-118. [Google Scholar]
 8. Walke SC, Chandrasekaran V, Mayya SS. Caregiver Burden among Caregivers of Mentally Ill Individuals and Their Coping Mechanisms. *J Neurosci Rural Pract.* 2018 Apr-Jun;9(2):180-185. doi: 10.4103/jnrp.jnrp_312_17. PMID: 29725166; PMCID: PMC5912021.
 9. Sultana Yesmin, Madhusri Manna, Kasturi Mandal, KathikaPattanayak, Sritama Nath Chowdhury. Assessment of family burden and it's co-relates among the caregivers of mentally ill patients in a OPD of selected mental hospital, West Bengal. *Int J Adv Psychiatric Nurs* 2022;4(1):26-34. DOI: 10.33545/26641348.2022.v4.i1a.78
 10. Sartorius N, Leff J, López-Ibor JJ, Maj M, Okasha A, World Psychiatric Association, editors. *Families and mental disorders from burden to empowerment.* Hoboken: Wiley; 2005. [Google Scholar]
 11. Steel Z, Marnane C, Iranpour C, Chey T, Jackson JW, Patel V, et al. The global prevalence of common mental disorders: a systematic review and meta-analysis 1980-2013. *Int J Epidemiol.* 2014;43(2):476-93. doi: 10.1093/ije/dyu038. [DOI] [PMC free article] [PubMed] [Google Scholar]
 12. Global Burden of Disease Study 2013 Collaborators. . Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: A systematic analysis for the Global Burden of Disease Study 2013 *Lancet.* 2015;386:743-800
 13. NIMHANS. . National Mental Health Survey of India, 2015-2016 Prevalence, Patterns and Outcomes, Supported by Ministry of Health and Family Welfare, Government of India, and Implemented by National Institute of Mental Health and Neurosciences (NIMHANS) 2016 Bengaluru NIMHANS
Available from: [http://www.nimhans.ac.in/sites/default/files/u197/NMHSReport\(Prevalencepatternsandoutcomes\)1](http://www.nimhans.ac.in/sites/default/files/u197/NMHSReport(Prevalencepatternsandoutcomes)1). [Last accessed on 2017 Nov 05]
 14. Katschnig H. Quality of life in mental disorders: Challenges for research and clinical practice *World Psychiatry.* 2006;5:139-45
 15. Dani MM, Thienhaus OJ. Characteristics of patients with schizophrenia in two cities in the U.S. and India *Psychiatr Serv.* 1996;47:300-1
 16. Thara R, Kamath S, Kumar S. Women with schizophrenia and broken marriages-doubly disadvantaged? Part II: family perspective *Int J Soc Psychiatry.* 2003;49:233-40
 17. Kaushik P, Bhatia MS. Burden and quality of life in spouses of patients with schizophrenia and bipolar disorder *Delhi Psychiatry J.* 2013;16:83-9
 18. Ofovwe CE, Osasona SO. Burden of caregiving among caregivers of patients with severe mental illnesses in Benin City, Nigeria. *Afr Health Sci.* 2022 Jun;22(2):657-667. doi: 10.4314/ahs.v22i2.75. PMID: 36407400; PMCID: PMC9652689.
 19. Cham CQ, Ibrahim N, Siau CS, Kalaman CR, Ho MC, Yahya AN, Visvalingam U, Roslan S, Abd Rahman FN, Lee KW. Caregiver Burden among Caregivers of Patients with Mental Illness: A Systematic Review and Meta-Analysis. *Healthcare (Basel).* 2022 Nov 30;10(12):2423. doi: 10.3390/healthcare10122423. PMID: 36553947; PMCID: PMC9777672.
 20. Chadda, R.K., Singh, T.B. & Ganguly, K.K. Caregiver burden and coping. *Soc Psychiat Epidemiol* 42, 923-930 (2007). <https://doi.org/10.1007/s00127-007-0242-8>
 21. von Kardorff, E., Soltaninejad, A., Kamali, M., & Eslami Shahrabaki, M. (2015). Family caregiver burden in mental illnesses: The case of affective disorders and schizophrenia - a qualitative exploratory study. *Nordic Journal of Psychiatry*, 70(4), 248-254. <https://doi.org/10.3109/08039488.2015.1084372>
 22. Savera Chothe. A Study to Assess the Level of Burden among the Caregivers of Mentally Ill Patients in Selected Hospitals of Imphal (Manipur). *The Nursing Journal of India.* May - June 2023 | Vol. CXIV, No. 3
 23. Gideon, Jayant & Ranjini, J & v, Richa. (2023). A Study to assess the burden of care and Coping strategies among caregivers of mentally ill patients in selected hospital, Dehradun, Uttarakhand. *International Journal of*

- Nursing Education and Research. 2. 293-298. 10.52711/2454-2660.2023.00066.
24. Mr. Nandakumar S. A Study to Assess the Level of Family Burden among the Caregivers of Schizophrenic Patients at a Mental Health Center in Salem. *RGUHS Journal of Nursing Sciences*. Volume: 11, Issue: 1, Year:2021, Page no. 11-13, DOI:10.26715/rjns.11_1_4
 25. Ayalew M, Workicho A, Tesfaye E, Hailesilasie H, Abera M. Burden among caregivers of people with mental illness at Jimma University Medical Center, Southwest Ethiopia: a cross-sectional study. *Ann Gen Psychiatry*. 2019 Jun 24;18:10. doi: 10.1186/s12991-019-0233-7. PMID: 31285750; PMCID: PMC6591984.
 26. Bhandari, Anup &Marahatta, Kedar & Rana, Mita & Ojha, S.P. & Regmi, M.P.. (2017). Caregiving Burden Among Family Members Of People With Mental Illness. *Journal of Psychiatrists' Association of Nepal*. 4. 36. 10.3126/jpan.v4i1.16741.
 27. C.J.Maikandaan,UmeshG,Vivian Kapil V,ZubaidaSultana.S Caregiver burden in family members of patients with mental illness *Global Journal For Research Analysis*, Vol.5,Issue :6 June 2016
 28. Gangadhar Naik¹, Vaishnavi Kommisetty², B.Renuka³, Pujari Ravikumar⁴, S. Pushpalatha⁵. Assessment of caregiver burden in psychiatric illness patients admitted in tertiary care hospital. *International Journal of Academic Medicine and Pharmacy* (www.academicmed.org) ISSN (O): 2687-5365; ISSN (P): 2753-6556
 29. Dharni J, Tuladhar S. Burden of Care among Caregivers of Mentally Ill Patients. *J Pharm Pract Community Med*. 2021;7(3):44-5.
 30. S., Sachin & Suresh, Venkatesan & H.N, Ravindra. (2014). A descriptive study to assess the burden among family care givers of mentally ill clients. *IOSR Journal of Nursing and Health Science*. 3. 61-67. 10.9790/1959-03346167.
 31. Bhandari, Anup &Marahatta, Kedar & Rana, Mita & Ojha, S.P. & Regmi, M.P.. (2017). Caregiving Burden Among Family Members Of People With Mental Illness. *Journal of Psychiatrists' Association of Nepal*. 4. 36. 10.3126/jpan.v4i1.16741.
 32. Mihan R, Mousavi SB, Khodaie Ardakani MR, Rezaei H, Hosseinzadeh S, Nazeri Astaneh A, Alikhani R. Comparison of Caregivers' Burden among Family Members of Patients with Severe Mental Disorders and Patients with Substance Use Disorder. *Iran J Psychiatry*. 2023 Apr;18(2):183-190. doi: 10.18502/ijps.v18i2.12369. PMID: 37383957; PMCID: PMC10293690.
 33. Ayalew, M., Workicho, A., Tesfaye, E. et al. Burden among caregivers of people with mental illness at Jimma University Medical Center, Southwest Ethiopia: a cross-sectional study. *Ann Gen Psychiatry* 18, 10 (2019). <https://doi.org/10.1186/s12991-019-0233-7>
 34. Nenobais, Aplonia&Jatimi, Atika &Jufriyanto, Muh. (2019). Family Burden for the Caregivers of People with Mental Disorders: A Systematic Review. *Jurnal Ners*. 14. 26. 10.20473/jn.v14i3.16971.
 35. Vir Vikram Sahdev Singh, Dr. R. Premavathy. Assess the Caregivers Burden, Stress and Coping Mechanism among Family Caregivers of Schizophrenic Patient - A Systematic Review. *International Journal of Creative Research Thoughts (IJCRT)*. Volume 10, Issue 3 March 2022 | ISSN: 2320-2882
 36. Thakur, Varsha; Nagarajan, Padmavathi¹; Rajkumar, Ravi Philip². Coping and Burden among Caregivers of Patients with Major Mental Illness. *Indian Journal of Social Psychiatry* 38(1):p 63-68, Jan-Mar 2022. | DOI: 10.4103/ijsp.ijsp_160_20
 37. Cheng, WL., Chang, CC., Griffiths, M.D. et al. Quality of life and care burden among family caregivers of people with severe mental illness: mediating effects of self-esteem and psychological distress. *BMC Psychiatry* 22, 672 (2022). <https://doi.org/10.1186/s12888-022-04289-0>
 38. Ebrahim, O.S., Al-Attar, G.S.T., Gabra, R.H. et al. Stigma and burden of mental illness and their correlates among family caregivers of mentally ill patients. *J. Egypt. Public. Health. Assoc.* 95, 31 (2020). <https://doi.org/10.1186/s42506-020-00059-6>
 39. C Samele, N Manning,Level of caregiver burden among relatives of the mentally ill in South Verona,Europe, *Psychiatry*,Volume 15, Issue 3,2000,Pages 196-204,ISSN 0924-9338,
 40. Udoh EE, Omorere DE, Sunday O, Osasu OS, Amoo BA (2021) Psychological distress and burden of care among family

- caregivers of patients with mental illness in a neuropsychiatric outpatient clinic in Nigeria. *PLoS ONE* 16(5): e0250309. <https://doi.org/10.1371/journal.pone.0250309>
41. Richa, Mano Ranjini J, Gideon J. A Study to assess the burden of care and Coping strategies among caregivers of mentally ill patients in selected hospital, Dehradun, Uttarakhand. *International Journal of Nursing Education and Research*. 2023; 11(4):293-8. doi: 10.52711/2454-2660.2023.00066 Available on: <https://ijneronline.com/AbstractView.aspx?PID=2023-11-4-5>
 42. Gupta P, Bharti P, Bathla M, Singh AH, Bhusri L. A cross-sectional study to assess the caregiver burden and the quality of life of caregivers of the patients suffering with psychiatric illness. *Ind Psychiatry J*. 2022 Jan-Jun;31(1):151-157. doi: 10.4103/ipj.ipj_228_20. Epub 2022 May 20. PMID: 35800861; PMCID: PMC9255620.
 43. Vaishali Tendolkar. "A Study To Assess The Level Of Stress Among Caregivers Of Individuals With Mental Disorders Attending Psychiatric Opd Of Selected Hospitals In The City." *European Journal of Molecular & Clinical Medicine* (2021): n. pag. Print.
 44. Rane, Nayna P., et al. "A Study To Assess The Level Of Stress Among Caregivers Of Individuals With Mental Disorders Attending Psychiatric Opd Of Selected Hospitals In The City." *European Journal of Molecular and Clinical Medicine*, vol. 8, no. 1, Wntr 2021, pp. 382+. *GaleAcademicOneFile*, link.gale.com/apps/doc/A698747675/AONE?u=anon-713cd723&sid=googleScholar&xid=5a97a6f8. Accessed 11 Feb. 2025.
 45. Souza, A.L.R., Guimarães, R.A., de Araújo Vilela, D. et al. Factors associated with the burden of family caregivers of patients with mental disorders: a cross-sectional study. *BMC Psychiatry* 17, 353 (2017). <https://doi.org/10.1186/s12888-017-1501-1>
 46. Rahmani F, Roshangar F, Gholizadeh L, Asghari E. Caregiver burden and the associated factors in the family caregivers of patients with schizophrenia. *Nurs Open*. 2022 Jul;9(4):1995-2002. doi: 10.1002/nop2.1205. Epub 2022 Mar 28. PMID: 35347867; PMCID: PMC9190687.
 47. Sultana Yesmin, Madhusri Manna, Kasturi Mandal, Kathika Pattanayak and Sritama Nath Chowdhury. "Assessment of Family Burden and It's Co-Relates among the Caregivers of Mentally Ill Patients in a OPD of Selected Mental Hospital, West Bengal." *International Journal of Advanced Psychiatric Nursing* 2022; 4(1): 26-34 <http://www.psychiatricjournal.net/> DOI:<https://doi.org/10.33545/26641348>. 2022.v4.i1a.78