

## Osteoma of left Temporal Bone - A Rare Case Report

Dr Gurumani Sriraman, Professor and HOD, Department of Otorhinolaryngology, Andaman & Nicobar Islands Institute of Medical Sciences, Andaman and Nicobar Islands 744104.

Email: [sriguruvenkatesh2014@gmail.com](mailto:sriguruvenkatesh2014@gmail.com)

Dr Deepthi E, Senior Resident, Department of Otorhinolaryngology, Andaman & Nicobar Islands Institute of Medical Sciences, Andaman and Nicobar Islands 744104.

*Received Date: 03/06/2025*

*Revised Date: 11/07/2025*

*Accepted Date: 14/08/2025*

### Abstract

**Background:** Osteoma is a benign, slow-growing tumour that develops from mature lamellar bone tissue. It is a single pedunculate mass which often happens unilaterally. The external auditory canal is more frequently affected by osteomas than other parts of the temporal bone. The purpose of this case report is to review this rare mastoid osteoma and discuss differential diagnosis and treatment. **Case Report:** 17-year-old female patient came to the outpatient clinic with a progressively enlarging postauricular mass for a duration of 10 years. The examination revealed that there was a mass of 2.0 x 1.2 cm on the left mastoid region that was hard and painless. HRCT temporal bone indicated Solitary osteoma observed from the outer table of the mastoid part of the temporal bone with a soft swelling of the adjacent soft tissues. **Conclusion:** This is a tumour of the temporal bone over the mastoid area that occurs rarely. The HRCT temporal bone eliminates major differentials and provides guidance in managing osteomas by visualizing tumour extension. Surgery treatment is required and should be considered depending on the cosmetic deformity and symptoms.

**Keywords:** Osteoma, left Temporal Bone.

### Introduction

Temporal bone osteoma is uncommon primary benign tumour of the temporal bone[1]. Those located in the mastoid and squamous parts of temporal bone may cause cosmetic deformity such as mastoid mass or an auricular protrusion [2]. In the skull they are commonly located in the fronto-ethmoid region. In differential diagnosis, other neoplasms of mastoid region including exostosis, osteosarcoma and osteoblastic metastasis should be considered [3]. The tumour has tendency to occur in all the bones of the body, however its occurrence in the temporal bone and middle ear is extremely rare.

### Case Report

17-year-old female patient came to the outpatient clinic with a progressively enlarging postauricular mass for a duration of 10 years (Fig. 1). There was no accompanying symptom of earache, discharge, hearing loss, vertigo, no facial weakness or tinnitus. Examination revealed a 2.0 × 1.2 cm mass on the left mastoid region which was hard and painless. There was no induration or erythema on the overlying skin. Bilateral tympanic membranes were normal. HRCT temporal bone indicated Solitary osteoma seen arising from outer table of mastoid part of temporal bone with adjacent mild soft tissue swelling (Fig. 2). Middle and inner ear were normal and the



Fig. 1 Showing the postauricular osteoma.




Fig. 2 CT scan showing the osseous outgrowth over the left mastoid.

Facial nerve was not involved. Patient was posted for local excision of the mass under general anesthesia. A postauricular skin incision and periosteal elevation were performed. The mass was exposed under the periosteum (Fig. 3) and removed entirely by using a chisel. The base of the lesion was drilled with a diamond burr under continuous irrigation until the base of the mass is exposed (Fig. 4). Wound was closed in layers. Postoperative period was uneventful. The specimen was sent for histopathological examination which was consistent with osteoma. In the histologic examination varying degrees of osteoblastic and osteoclastic activities showing lamellar bone with fibrovascular bundles were noted (Hematoxylin and Eosin; original magnification,  $\times 10$ ).

### Discussion

Head & neck osteomas are very rare. Osteomas located in all portions of the temporal bone, including external auditory canal (the most common site), squama, mastoid, middle ear, glenoid fossa, eustachian tube, internal auditory canal, petrous apex and styloid process have been reported [2] [3] [5] -[9]. Mastoid portion is the second most common area for the existence of temporal bone osteomas following the external auditory canal [2].

Histologically there are three types of osteoma of the mastoid: osteoma compactum, osteoma cancellare, and osteoma cartilagineum. Clinically it is difficult to distinguish the type of osteoma because of the similarity of the symptoms and objective signs [10]. Compact osteomas have a wider base and are very slow growing whereas spongy osteomas are more likely to be pedunculated and grow relatively faster.

Trauma, surgery, radiotherapy, chronic infection and pituitary dysfunction are all included in the etiology of mastoid osteomas. However the precise etiology is still unknown and considered to be a true bone tumor [11] - [14].

At the time of presentation the size of osteoma is generally not bigger than 3 cm [2] [18]. Superficial osteomas have smooth surface, bony hard in consistency and the overlying skin is not involved [2]. The presented lesion caused cosmetic deformity as a result of gradual growth to the size of approximately  $4 \times 3$  cm over 10 years. Clinically mastoid osteomas rarely cause pain or inflammation. Pain occurs when the growth of the lesion involves the inner table of the temporal bone [19]. Pain may be localized in the ear region, the tympanic membrane or in the neck. The latter can be due to the irritation of the greater auricular or small occipital nerves [10]



Fig. 3 Osteoma on elevation of the periosteum.



Fig. 4 View of the base of the osteoma.

Osteomas can cause functional symptoms. The neoplasm may infiltrate the cortex, causing posterior wall to be pushed forward, resulting in obstruction of the external auditory canal and conductive hearing loss.

Osteomas can arise from the inner table of squamous and petrous part of the temporal bone resulting in brain compression and generalized epileptic seizure [20] .

Pressure-induced pain can be referred to the neck, temporomandibular joint or middle ear. In the presence of external ear canal obstruction, conductive hearing loss and chronic suppuration may occur [21] [22] . A mastoid osteoma compressing the posterior fossa structures and causing intracranial complications has been reported by Van Dellen [23] .

In differential diagnosis of mastoid osteomas, osteosarcoma; osteoblastic metastasis; isolated eosinophilic granuloma; Paget's disease; giant cell tumor; osteoid osteoma; calcified meningioma and monostotic fibrous dysplasia should be considered. Radiologic borders of these lesions are less clear than those of osteomas [3] . HRCT scan demonstrates an osteoma as well demarcated, dense, sclerotic outgrowth from the mastoid bone. Surgery is indicated for both cosmetic reasons and confirmation of the diagnosis of osteomas of the mastoid and squamous portion of the temporal bone. Exostosis and osteoma are very similar in most aspects however exostosis lacks fibrovascular channels [24] . Also osteomas are bony growths that are single, unilateral and pedunculated and mostly arise from tympanomastoid or tympanosquamous suture lines laterally [6] whereas exostoses are multiple, usually bilateral and broad based and are found medial to sutures of the temporal bone [11] .

The osteoma must completely be excised until normal mastoid air cells are exposed. These lesions are limited to the external cortex generally hence a cleavage plane is encountered where the osteoma meets the normal bone. In mastoid osteomas extending into the bony labyrinth and facial nerve, complete removal might not be indicated since the damage to these structures is likely. Recurrence is rare [2] [21] .

Tumours involving middle ear and inner ear are small and tend to remain small so observation is preferred unless symptomatic. When there is functional disorder like hearing loss or vertigo judicious surgical removal maybe undertaken. Surgical management of internal auditory canal osteoma is varied. Both middle cranial fossa approach and suboccipital approach have been employed [25] . Approach generally depends on location, size, preference and experience of the surgeon.

Gardner's syndrome must be kept in mind in cases of multiple osteomas. Gardner's syndrome is characterized by multiple intestinal polyps, epidermoid inclusion cysts, fibromas of the skin

and mesentery and osteomas. Osteomas in Gardner's syndrome have a predilection for membranous bones and the mandible and maxilla are more commonly involved [26].

### **Conclusion**

Osteoma of temporal bone over mastoid area is a very rare tumour. HRCT temporal bone eliminates the main differentials and guides in management of osteomas by visualising tumour extension. Treatment is surgical and must be considered according to the cosmetic deformity and the symptoms.

### **References**

1. Unal, O.F., Tosun, F., Yetiser, S. and Dundar, A. (2000) Osteoma of the Middle Ear. *International Journal of Pediatric Otorhinolaryngology*, 52, 193-195. [http://dx.doi.org/10.1016/S0165-5876\(00\)00286-X](http://dx.doi.org/10.1016/S0165-5876(00)00286-X) [Citation Time(s):1]
2. Denia, A., Perez, F., Canalis, R.R., et al. (1979) Extracanalicular Osteomas of the Temporal Bone. *ACTA Oto-Laryngologica*, 105, 706-709. <http://dx.doi.org/10.1001/archotol.1979.00790240020005> [Citation Time(s):6]
3. Estrem, S.A., Vessely, M.B. and Oro, J.J. (1993) Osteoma of the Internal Auditory Canal. *Otolaryngology, Head and Neck Surgery*, 108, 293-297 [Citation Time(s):3]
4. Pérez, á.D.D., Romero, R.R., Durán, D.E., Montaña, P.R., Bernal, R.A. and Rodríguez, C.M. (2011) El osteoma en la mastoides, ¿un hallazgo incidental? *Acta Otorrinolaringológica Española*, 62, 140-143. <http://dx.doi.org/10.1001/archotol.1940.00660010852006> [Citation Time(s):1]
5. Burton, D.M. and Gonzalez, C. (1991) Mastoid Osteomas. *Ear, Nose Throat Journal*, 70, 161-162 [Citation Time(s):1]
6. Yamasoba, T., Harada, T., Okuno, T., et al. (1990) Osteoma of the Middle Ear. Report of a Case. *Archives of Otolaryngology—Head and Neck Surgery*, 116, 1214-1216. <http://dx.doi.org/10.1001/archotol.1990.01870100108025> [Citation Time(s):1]
7. D'Ottovai, L.R., Piccirillo, E., De Sanctis, S., et al. (1997) Mastoid Osteomas: Review of the Literature and Presentation of Two Clinical Cases. *Acta Otorhinolaryngologica Italica*, 17, 136-139
8. Gungor, A., Cincik, H., Poyrazoglu, E., Saglam, O. and Candan, H. (2004) Mastoid Osteomas: Report of Two Cases. *Otology Neurotology*, 25, 95-97. <http://dx.doi.org/10.1097/00129492-200403000-00002>
9. Birrell, J.F. (1978) Osteoma of the Mastoid. *Journal of the Royal College of Surgeons of Edinburgh*, 23, 305-309 [Citation Time(s):1]
10. Cinelli, A.A. (1941) Osteoma Eburneum of the Mastoid. *Archives of Otolaryngology, Head and Neck Surgery*, 33, 421-424. <http://dx.doi.org/10.1001/archotol.1941.00660030425008> [Citation Time(s):2]
11. Fenton, J.E., Turner, J. and Fagan, P.A. (1996) A Histopathologic Review of Temporal Bone Exostoses and Osteomata. *Laryngoscope*, 106, 624-628. <http://dx.doi.org/10.1097/00005537-199605000-00020> [Citation Time(s):2]
12. Graham, M.D. (1979) Osteomas and Exostoses of the External Auditory Canal. *Journal of Otolaryngology*, 88, 566-572
13. Tran, L.P., Grungfast, K.M. and Selesnick, S.H. (1996) Benign Lesions of the External Auditory Canal. *Otolaryngologic Clinics of North America*, 29, 807-825
14. Fisher, E.W. and McManus, T.C. (1994) Surgery of the External Auditory Canal Exostoses and Osteomata. *The Journal of Laryngology & Otology*, 108, 106-110. <http://dx.doi.org/10.1017/S0022215100126027> [Citation Time(s):1]

15. Friedberg, S.A. (1938) Osteoma of Mastoid Process. Archives of Otolaryngology, Head and Neck Surgery, 28, 20-26. <http://dx.doi.org/10.1001/archotol.1938.00650040027003> [Citation Time(s):2]
16. Thomas, R. (1964) Familial Osteoma of the Middle Ear. The Journal of Laryngology & Otology, 78, 805-807. <http://dx.doi.org/10.1017/S0022215100062794> [Citation Time(s):1]
17. Ohhashi, M., Terayama, Y. and Mitsui, H. (1984) Osteoma of the Temporal Bone: A Case Report. Nippon Jibiinkoka Gakkai Kaiho, 87, 590-595. <http://dx.doi.org/10.3950/jibiinkoka.87.590> [Citation Time(s):1]
18. Guerin, N., Chauveau, E., Julien, M., Dumont, J.M. and Merignargues, G. (1996) Osteoma of the Mastoid Bone: Report of Two Cases. Annals of Otology, Rhinology, and Laryngology, 117, 127-132 [Citation Time(s):1]
19. Stuart, E.A. (1940) Osteoma of the Mastoid: Report of a Case with Investigations of the Constitutional Background. Archives of Otolaryngology, Head and Neck Surgery, 31, 838-854. <http://dx.doi.org/10.1001/archotol.1940.00660010852006> [Citation Time(s):1]
20. Bhandari, Y.S. and Jones, R.A.C. (1972) Osteoma of the Middle Cranial Fossa. Journal of Neurosurgery, 37, 610-612. <http://dx.doi.org/10.3171/jns.1972.37.5.0610> [Citation Time(s):1]
21. Probst, L.E., Shankar, L. and Fox, R. (1991) Osteoma of the Mastoid Bone. Journal of Otolaryngology, 20, 228-230 [Citation Time(s):2]
22. Singh, I., Sanasam, J.C., Bhatia, P.L. and Singh, L.S. (1979) Giant Osteoma of the Mastoid. Ear, Nose and Throat Disorders, 58, 243-245 [Citation Time(s):1]
23. Van Dellen, J.R. (1977) A Mastoid Osteoma Causing Intracranial Complications: A Case Report. The South African Medical Journal, 51, 597-598 [Citation Time(s):1]
24. Sheehy, J.J. (1982) Diffuse Exostoses and Osteomata of the External Auditory Canal: A Report of 100 Cases. Otolaryngology, Head and Neck Surgery, 90, 337-342 [Citation Time(s):1]
25. Clerico, D.M., Jahn, A.F. and Fontanella, S. (1994) Osteoma of the Internal auditory Canal. Case Report and Literature Review. Annals of Otology, Rhinology, and Laryngology, 103, 9-23 [Citation Time(s):1]
26. Smud, D., Augustin, G., Kekez, T., Kinda, E., Majerovic, M. and Jelincic, Z. (2007) Gardner's Syndrome: Genetic Testing and Colonoscopy Are Indicated in Adolescents and Young Adults with Cranial Osteomas: A Case Report. World Journal of Gastroenterology, 13, 3900-3903 [Citation Time(s):1].